

# Key access and utilization descriptors for chickpea genetic resources

This list consists of an initial set of characterization and evaluation descriptors for chickpea (*Cicer arietinum* L.) genetic resources utilization. This strategic set of descriptors, together with passport data, will become the basis for the global accession level information portal being developed by Bioversity International with the financial support of the Global Crop Diversity Trust (GCDT). It will facilitate access to and utilization of chickpea accessions held in genebanks and does not preclude the addition of further descriptors, should data subsequently become available.

Based on the comprehensive list 'Descriptors for Chickpea (*Cicer arietinum* L.)' published by ICRISAT, ICARDA and IBPGR (now Bioversity International) in 1993, the list builds on the results of the Global Public Goods Activity 4.2.1.1, particularly with regards to those descriptors highlighted as the most important diagnostic and breeding traits. It was subsequently compared and harmonized with a number of sources such as UPOV technical guidelines for Chick-Pea (2005), 'Descriptors for CHICKPEA' (USDA, ARS, GRIN), 'Core Collection of Chickpea as a Means to Enhance Utilization of Genetic Resources in Crop Improvement' (ICRISAT-website), 'Global Strategy for the *Ex situ* Conservation of Chickpea (*Cicer* L.)' (GCDT, 2008), as well as with those descriptors that were awarded funds for further research by the GCDT in 2008 Evaluation Awards Scheme (EAS). This list was further refined during a meeting held at the National Bureau of Plant Genetic Resources (NBPGR, India) in June 2009. Several scientists from NBPGR and the Indian Agricultural Research Institute (IARI) participated.

A worldwide distribution of experts was involved in an online survey to define a first priority set of descriptors to describe, to access and to utilize chickpea genetic resources. This key set was afterwards validated by a Core Advisory Group (see 'Contributors') led by Dr M. Imtiaz of the International Center for Agricultural Research in the Dry Areas (ICARDA), Dr M.C. Kharkwal (IARI) and Dr Hari D. Upadhyaya of the International Crops Research Institute for the Semi-Arid Tropics (ICRISAT).

Biotic and abiotic stresses included in the list were chosen because of their wide geographic occurrence and significant economic impact at a global level.

Numbers in parentheses on the right-hand side are the corresponding descriptor numbers listed in the 1993 publication. Descriptors with numbers ending in 'letters' are either modified or are new descriptors that were added during the development of the list below.

## PLANT DATA

### Stem/foilage pigmentation

(4.1.1)

Observed before flowering. Indicate whether the pigmentation is on stems or leaves in the descriptor **Notes**

- 1 No anthocyanin (light green)
- 3 No anthocyanin (green)
- 5 Low anthocyanin (partly light purple)
- 7 High anthocyanin (predominantly purple)
- 9 Highly purple

### **Days to 50% flowering** (4.2.1)

Number of days from sowing (or first rain sufficient for germination under rainfed conditions) until 50% of the plants have started to flower

### **Days to maturity** (4.2.2)

Number of days from sowing (or first rain sufficient for germination under rainfed conditions) until 90% of the pods have matured and turned yellow

### **Number of seeds per pod** (4.2.3)

Average number of 10 pods each from five representative plants. At maturity

### **Flower colour** (4.2.4)

In most cases pink and blue flowers have veins of a darker shade in the flag, while the tip of the keel is also darker. The classes are ranges rather than only the shades of the reference colours. Royal Horticultural Society (RHS) colour codes are given in parentheses beside descriptor states

- 1 Blue (violet-blue group 97B)
- 2 Light blue (violet-blue group 97C)
- 3 Dark pink (red-purple group 64D)
- 4 Pink (red-purple group 63D)
- 5 Light pink (red-purple group 69C)
- 6 White (white group 155D)
- 7 White-pink striped (white group 155D, red-purple group 63D)

### **Number of pods per plant** (4.2.8)

Average number of pods taken from five representative plants. At maturity

### **Seed shape** (4.3.1)

- 1 Angular, ram's head (most desi cultivars)
- 2 Irregular rounded, owl's head (most kabuli cultivars)
- 3 Pea-shaped, smooth round (intermediate types)

### **Seed testa texture** (4.3.2)

- 1 Rough (pea-shaped)
- 2 Smooth
- 3 Tuberculated (sticky surface)

**Seed colour** (4.3.3)

Royal Horticultural Society (RHS) colour codes are given in parentheses beside descriptor states

- 1 Black (black group 202A, 202B; brown group 200A)
- 2 Brown (greyed-orange group 177B)
- 3 Light brown (greyed-orange group 177C)
- 4 Dark brown (greyed-orange group 177A)
- 5 Reddish brown (greyed-orange group 166C)
- 6 Greyish brown (brown group 200D)
- 7 Salmon brown (greyed-orange group 165C)
- 8 Grey (greyed-green group 196A)
- 9 Brown beige (greyed-orange group 173D)
- 10 Beige (greyed-orange group 165D)
- 11 Yellow (greyed-orange group 164B)
- 12 Light yellow (greyed-orange group 164C)
- 13 Yellow brown (greyed-orange group 165C)
- 14 Orange yellow (greyed-orange group 168D)
- 15 Orange (greyed-orange group 168C)
- 16 Yellow beige (orange-white group 159C)
- 17 Ivory white (orange-white group 159C)
- 18 Green (greyed-green group 191A; grey group 201A; greyed-orange group 166B)
- 19 Light green (greyed-green group 193B)
- 20 Variegated
- 21 Black brown mosaic (black group 202A; greyed-orange group 177E)

**100-seed weight [g]** (4.3.5)

Measured at 10% (air-dry) moisture content

**Growth habit** (6.1.1)

The angle of the branches from the vertical axis at the pod filling stage

- 1 Prostrate (branches flat on the ground,  $>80^\circ$ )
- 2 Spreading ( $61-80^\circ$  from vertical)
- 3 Semi-spreading ( $26-60^\circ$  from vertical)
- 4 Semi-erect ( $16-25^\circ$  from vertical)
- 5 Erect ( $0-15^\circ$  from vertical)

**Number of primary branches** (6.1.4.1)

Average number of basal primary branches per plant taken from five representative plants

**Plant canopy height [cm]** (6.1.5)

Average canopy height of five representative plants. Recorded at maturity

**Seed yield per plant [kg ha<sup>-1</sup>]** (6.2.2.2)**Seed protein content [% DW]** (6.3.1.1)

Whole seed crude protein using the dye-binding method or automatic protein analyzer

**ABIOTIC STRESSES**

**Reaction to drought** (7.5)

**Reaction to salinity** (7.X)

**BIOTIC STRESSES**

**Ascochyta blight** (*Ascochyta rabiei*) (8.1.2)

**Grey mould** (*Botrytis cinerea*) (8.1.3)

**Fusarium wilt** (*Fusarium oxysporum* f. sp. *ciceri*) (8.1.4)

**Pod borer** (*Helicoverpa armigera*) (8.6.1)

**NOTES**

Any additional information may be specified here, particularly that referring to the category '99=Other' present in some of the descriptors above.

## CONTRIBUTORS

Bioversity is grateful to all the scientists and researchers who have contributed to the development of this strategic set of 'Key access and utilization descriptors for chickpea genetic resources', and in particular to Dr M. Imtiaz, Dr M.C. Kharkwal and Dr Hari Upadhyaya for providing valuable scientific direction. Adriana Alercia provided technical expertise and guided the entire production process.

## CORE ADVISORY GROUP

**Muhammad Imtiaz**, International Center for Agricultural Research in the Dry Areas (ICARDA), Syria

**M.C. Kharkwal**, Division of Genetics, Indian Agricultural Research Institute (IARI), India

**Hari D. Upadhyaya**, International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), India

C. Bharadwaj, Division of Genetics, Indian Agricultural Research Institute (IARI), India

François Boulineau, Groupe d'Etude et de contrôle des Variétés et des Semences (GEVES), France

Clarice J. Coyne, United States Department of Agriculture, Agricultural Research Service (USDA, ARS), USA

Paola Crinò, Agenzia nazionale per le nuove tecnologie, l'energia e lo sviluppo economico sostenibile (ENEA), Italy

R.P. Dua, Indian Council of Agricultural Research (ICAR), India

Matthias Kotter, Leibniz Institute of Plant Genetics and Crop Plant Research (IPK), Germany

Ulrike Lohwasser, Leibniz Institute of Plant Genetics and Crop Plant Research (IPK) Department of Genebank, Research Group Resources Genetics and Reproduction, Germany

S.R. Pandravada, National Bureau of Plant Genetic Resources (NBPGR), Regional Station, Hyderabad, India

## REVIEWERS

### Australia

Tanveer Khan, Department of Agriculture and Food

Bob Redden, Department of Primary Industries Victoria

Kadambot Siddique, The University of Western Australia

### Bangladesh

Mamtazul Haque, Bangladesh Agricultural Research Institute

### Canada

Axel Diederichsen, Plant Gene Resources of Canada, Agriculture and Agri-Food Canada

Bunyamin Taran, Crop Development Centre, University of Saskatchewan

### Cuba

Tomás Shagarodsky Scull, Instituto de Investigaciones Fundamentales en la Agricultura Tropical (INIFAT)

### **India**

C.L.L. Gowda, International Crops Research Institute for the Semi-Arid Tropics (ICRISAT)

Jitendra Kumar, Indian Agricultural Research Institute (IARI)

S.K. Mishra, National Bureau of Plant Genetic Resources (NBPGR)

M. Thimma Reddy, International Crops Research Institute for the Semi-Arid Tropics (ICRISAT)

Mamta Sharma, International Crops Research Institute for the Semi-Arid Tropics (ICRISAT)

Shivali Sharma, International Crops Research Institute for the Semi-Arid Tropics (ICRISAT)

Sube Singh, International Crops Research Institute for the Semi-Arid Tropics (ICRISAT)

Shyam Singh Yadav, Indian Agricultural Research Institute (IARI)

### **Israel**

Shahal Abbo, Hebrew University of Jerusalem

### **Mexico**

José Antonio Garzón-Tiznado, Universidad Autónoma De Sinaloa

### **Pakistan**

Ahmad Zahoor, National Agricultural Research Centre (NARC)

### **Slovak Republic**

Gabriela Antalíková, Plant Production Research Centre, Research Institute of Plant Production (PPRC, RIPP) Piešťany

### **Spain**

Rafael M. Jiménez-Díaz, University of Córdoba

### **The Netherlands**

L. J. G. van der Maesen, Wageningen University

### **USA**

Fred Muehlbauer, United States Department of Agriculture, Agricultural Research Service (USDA, ARS)